### Total vegetation cover soil protection Region:LGA Lockhart (A) NSW

This report describes vegetation protecting the soil surface from erosion during a chosen month compared to previous years. This report has been generated using MODIS fractional vegetation cover information available in Rangelands and Pasture Productivity (RAPP) map tool https://map.geo-rapp.org/#australia. The report is based on 500 metre pixel data on monthly time steps.

Land use forest cover:

**Date: May 2023** 

Results can be shown for the whole region (polygon), and separated by land use and forest cover classes which are likely to show different cover patterns and targets. Land use is divided into four broad classes: Conservation and natural environments, Agriculture, production native forests and plantation forests (no report), and other (no report). Agriculture is divided into grazing, crops and horticulture which are sub-divided into non-irrigated and irrigated. If forest is present land use is further divided into: non-forest, woodland forest and non-woodland forest. The area of each land use and forest class are shown as a map and chart. The report content is repeated for each land use and forest cover class that covers at least 1% of the area of the chosen region. Total vegetation Cover:

The total vegetation cover indicates where soil is likely to be protected from wind and or water hillslope erosion. Total vegetation cover for this month is shown on a map and chart classified into 4 classes.

- 71-100% High cover protected from wind and usually water erosion (high rainfall, steep slopes, and erodible soils may need greater than 80, 90, 95 and up to 100% cover)
  - 51-70% Moderate cover protected from wind erosion
  - 31-50% Low cover not protected
  - 0-30% Very Low cover not protected

Erosion protection: Wind erosion 50% total vegetation cover

The vegetation cover threshold required to prevent soil erosion is usually 50% to reduce wind erosion, 70% or 80% to reduce water (hillslope) erosion depending on the steepness and rainfall. Areas protected from erosion for the month:

- Map: water erosion protection (>70% cover) percentage area and hectares.
- Map: wind erosion protection (>50% cover) percentage area and hectares.

Comparison with previous years:

- Map: anomaly comparing this month to the average cover from the same month in previous years.
- Map: deciles rank of month against the same month in previous years.

Anomalies and deciles until September 2019 are calculated comparing to the same months 2001 to 2019. Extra monthly data will be used to calculate anomalies and deciles post September 2019 as they become available. Time series monthly from January 2001 to current:

#### **Erosion protection**

- Wind erosion protection time series: percentage of the area of the region with greater than 50% cover for each month (orange lines). Horizontal lines are 10th (cover target) and 50th percentiles.
- Water erosion protection time series: percentage of the area of the region with greater than 70% cover for each month (blue line). Horizontal lines are 10th (cover target) and 50th percentiles.

#### Rainfall

• Millimetres rainfall each month (black line).

Each time series is also stacked by year. The black line shows the current year of data.

Water erosion protection for higher rainfall and steeper slopes:

Water erosion protection on higher slopes. As slope increases, more cover is required to control water erosion. The thresholds reported are:

- the percentage area with pixels greater than 80% total cover.
- the percentage area with pixels greater than 90% total cover.
- the percentage area with pixels greater than 95% total cover.

#### **Acknowledgment of data:**

- 1. http://www.agriculture.gov.au/abares/aclump/land-use/alum-classification
- 2. http://www.agriculture.gov.au/abares/forestsaustralia/sofr/sofr-2018
- 3. https://www.dpi.nsw.gov.au/agriculture/pastures-and-rangelands/establishment-mgmt/production-management2/groundcover
- 4. MODIS Fractional cover algorithm:

https://doi.org/10.4225/08/5848a3f19a7b3











#### **Vegetation Cover May 2023**


#### Land use and forest cover




#### Proportion of each land class in area




#### Total Vegetation Cover [%]




Proportion of vegetation cover class in area




#### % Area protected from water erosion (>70%)



% Area protected from wind erosion (>50%)



#### Total Vegetation Cover Anomaly [%]



Deciles show where the pixel value lies in the record, from highest to lowest, for that month. That is, red pixels are in the lowest 10% of records for that month of the map using baseline from 2001 to 2019.


**Total Vegetation Cover Decile [%]** 











.


#### **Agriculture**

# Catchment Scale Land Use and Forests of Australia (2018) Derived from Catchment Scale Use of Australia (2018) Agriculture - Grazing - Non forest 2 Agriculture - Grazing - Non-woodland forest Use of Australia (2018) and Forests of Australia (2018)


# 80 - 92.6% 60 - 20 - 0.5 0.0 0.5 1.0 1.5 2.0 2.5 3.0 3.5 Land use class

**Proportion of each land class in area** 


Proportion of vegetation cover class in area



% Area protected from water erosion (>70%)



% Area protected from wind erosion (>50%)




Area not protected 0.0% of region (0 ha)


Area protected 100.0% of region (284,100 ha)

**Total Vegetation Cover Anomaly [%]** 

Total Vegetation Cover Decile [%]



Deciles show where the pixel value lies in the record, from highest to lowest, for that month. That is, red pixels are in the lowest 10% of records for that month of the map using baseline from 2001 to 2019.

















#### **Agriculture timeseries**













#### **Grazing**




#### **Proportion of each land class in area** 88.8% 80 60 20 8.3% -0.5 0.5 1.0 2.0 0.0 2.5 1.5 Land use class











#### % Area protected from water erosion (>70%)


% Area protected from wind erosion (>50%)






#### **Total Vegetation Cover Anomaly [%]**

**Total Vegetation Cover Decile [%]** 





Deciles show where the pixel value lies in the record, from highest to lowest, for that month. That is, red pixels are in the lowest 10% of records for that month of the map using baseline from 2001 to 2019.













#### **Grazing timeseries**











Water erosion historical monthly area protected (Total Veg Cov>70%)

#### **Grazing non forest**


#### Land use and forest cover




#### Total Vegetation Cover [%]



#### % Area protected from water erosion (>70%)




#### Total Vegetation Cover Anomaly [%]




Deciles show where the pixel value lies in the record, from highest to lowest, for that month. That is, red pixels are in the lowest 10% of records for that month of the map using baseline from 2001 to 2019.


#### **Proportion of vegetation cover class in area**



#### % Area protected from wind erosion (>50%)



#### Total Vegetation Cover Decile [%]















#### **Grazing non forest timeseries**











#### **Cropping**


#### Land use and forest cover




#### **Total Vegetation Cover [%]**



#### % Area protected from water erosion (>70%)




#### Total Vegetation Cover Anomaly [%]




Deciles show where the pixel value lies in the record, from highest to lowest, for that month. That is, red pixels are in the lowest 10% of records for that month of the map using baseline from 2001 to 2019.


#### Proportion of vegetation cover class in area



#### % Area protected from wind erosion (>50%)



#### Total Vegetation Cover Decile [%]















#### **Cropping timeseries**









## Lockhart\_(A) (289,400 ha and no data 305 ha) Percentage area and hectares protected with TVC threshold 30,50,70,80,90 and 95%

Land use and forest cover Class	area(ha)	above_30	above_50	above_70	above_80	above_90	above_95
Entire region	289,400	99.8% 288,850	99.7% 288,600	85.9% 248,525	57.2% 165,425	12.6% 36,375	2.4% 7,000
Agriculture	284,100	100.0% 284,100	99.9% 283,925	85.9% 244,150	57.0% 161,800	12.4% 35,175	2.3% 6,625
Grazing	21,125	100.0% 21,125	100.0% 21,125	97.6% 20,625	86.2% 18,200	33.7% 7,125	10.8% 2,275
Grazing non forest	18,750	100.0% 18,750	100.0% 18,750	97.3% 18,250	84.5% 15,850	32.8% 6,150	11.9% 2,225
Cropping	262,975	100.0% 262,975	99.9% 262,800	85.0% 223,525	54.6% 143,600	10.7% 28,050	1.7% 4,350







