This report provides information about vegetation covering the soil surface for a region during a single month with comparison to previous years. The total vegetation cover indicates where soil is likely to be protected from wind (>=50% total vegetation cover) and water/hillslope (>=70% total vegetation cover) erosion. Results are shown for the whole region (polygon) and also separated by land use and forest cover class. This is because different land use / forest cover classes are likely to have different cover patterns and targets.

[Karratha (C)]

The six maps and two graphs provide a report for the month with:

- Land use and forest cover information for the area:
 - o Map: Land use and forest cover
 - o Chart: Land use and forest cover area
- Total vegetation cover for this month:
 - o Map: total vegetation cover classified into 4 classes
 - o Chart: total vegetation cover percentage area classified into 4 classes
- Areas protected from erosion for the month:
 - o Map: water erosion protection (>70% cover) percentage area and hectares
 - o Map: wind erosion protection (>50% cover) percentage area and hectares
- Comparison with previous years:
 - o Map: anomaly compare this month to the average cover from the same month in previous years
 - o Map: deciles rank this month against the same month in previous years
- Time series from January 2001 to current:
 - o Wind erosion protection time series: percentage of the area of the region with greater than 50% cover for each month since January 2001 (orange line): Horizontal lines are 10th (cover target) and 50th percentiles. Vertical line is month of report.
 - o Water erosion protection time series: percentage of the area of the region with greater than 70% cover for each month since January 2001 (blue line): Horizontal lines are 10th (cover target) and 50th percentiles. Vertical line is month of report.
 - o Rainfall: millimetres rainfall each month (black line). Vertical line is month of report.
- Time series for each month stacked by year
 - o Same data as time series from January 2001 to current month, grouped by month. Black line is current year of data.
- Water erosion protection on higher slopes. As slope increases, more cover is required to control water erosion.

The thresholds reported are:

- o the percentage area with pixels greater than 80% total clover
- o the percentage area with pixels greater than 90% total clover
- o the percentage area with pixels greater than 95% total clover

The following pages repeat the above sequence for each land use and forest cover class. For example

- All agricultural lands, that is grazing, cropping plus Horticulture (depending on what land use is present)
- Grazing lands by forest classes if present
- Cropping lands
- Irrigation lands
- Protected areas by forest classes if present

The following pages repeat the above sequence for each land use and forest cover class if 1% or more of area makes up a land use and forest cover class. Four land uses are reported: Conservation and natural environments, Agriculture, production native forests and plantation forests, and other. Agriculture is further divided into grazing,

crops and horticulture are then divided into non-irrigated and irrigated. Land use is further divided by forest class if present: non-forest, woodland forest and non-woodland forest.

Explanatory notes:

This report has been generated using MODIS fractional vegetation cover information available in Rangelands and Pasture Productivity (RAPP) map tool. The report is based on an analysis of 500 metre pixel data on monthly time steps. Report uses baseline from January 2001 to September 2019 for each month to generate anomalies and deciles. Post September 2019 all similar months are used to calculate anomalies and deciles.

Acknowledgment of data:

- 1. http://www.agriculture.gov.au/abares/aclump/land-use/alum-classification
- 2. http://www.agriculture.gov.au/abares/forestsaustralia/sofr/sofr-2018
- 3. https://www.dpi.nsw.gov.au/agriculture/pastures-and-rangelands/establishment-mgmt/production-management2/groundcover
- 4. MODIS Fractional cover algorithm:

https://doi.org/10.4225/08/5848a3f19a7b3















Vegetation Cover Sep 2019


Proportion of each land class in area







Total Vegetation Cover [%]












% Area protected from water erosion (>70%)























Conservation and natural environments

Land use and forest cover




Total Vegetation Cover [%] Total Vegetation Cover [%] Total Vegetation Cover [%] Total Vegetation Cover [%]

50 - 51.3% 40 - 40.5% 30 - 51.3% 20 - 6.5%

Proportion of vegetation cover class in area

% Area protected from water erosion (>70%)






Total Vegetation Cover class


31%-50%


0-30%



51%-70%

71%-100%





















Conservation and natural environments timeseries










Conservation and natural environments non forest


Land use and forest cover




Total Vegetation Cover [%] Typic Toolo Typ


Proportion of vegetation cover class in area 50 - 51.5% 40 - 40.5% 30 - 20 - 10 - 6.3% Total Vegetation Cover class


% Area protected from water erosion (>70%)



























Conservation and natural environments non forest timeseries











Agriculture

Land use and forest cover




Total Vegetation Cover [%] Total Vegetation Cover [%] Trobattoolo T


Proportion of vegetation cover class in area





% Area protected from water erosion (>70%)



% Area protected from wind erosion (>50%)





















Agriculture timeseries



---- 2016

2017 --- 2018 --- 2019

Proportion of the area protected from water erosion (Total Vegetation Cover > 70%)




month

Ω


Grazing

Land use and forest cover




Total Vegetation Cover [%] Total Vegetation Cover [%] Trobattoolo T


Proportion of vegetation cover class in area





% Area protected from water erosion (>70%)



% Area protected from wind erosion (>50%)























Grazing timeseries






month






Grazing non forest

Land use and forest cover




Total Vegetation Cover [%] Teleradolo Te


Proportion of vegetation cover class in area





% Area protected from water erosion (>70%)



% Area protected from wind erosion (>50%)























Grazing non forest timeseries





month





Karratha_(C) (1,500,450 ha and no data 23,445 ha) Percentage area and hectares protected with TVC threshold 30,50,70,80,90 and 95%

Land use and forest cover Class	area(ha)	above_30	above_50	above_70	above_80	above_90	above_95
Entire region	1,500,450	98.4% 1,476,550	38.3% 574,925	2.2% 32,600	1.0% 15,200	0.4% 5,725	0.2% 2,850
Conservation and natural environments	471,141	98.3% 463,366	47.1% 221,709	6.5% 30,817	3.2% 15,063	1.2% 5,549	0.6% 2,736
Conservation and natural environments non forest	469,640	98.3% 461,862	46.9% 220,081	6.3% 29,707	3.1% 14,406	1.1% 5,347	0.6% 2,686
Agriculture	1,014,304	98.4% 998,543	34.3% 347,812	0.2% 1,871	0.0% 24	0.0% 24	0.0% 24
Grazing	1,014,304	98.4% 998,543	34.3% 347,795	0.2% 1,871	0.0% 24	0.0% 24	0.0% 24
Grazing non forest	1,014,304	98.4% 998,540	34.3% 347,691	0.2% 1,871	0.0% 24	0.0% 24	0.0% 24











