Total vegetation cover soil protection Region:NRM North West NSW NSW

Date: February 2006

This report describes vegetation protecting the soil surface from erosion during a chosen month compared to previous years. This report has been generated using MODIS fractional vegetation cover information available in Rangelands and Pasture Productivity (RAPP) map tool https://map.geo-rapp.org/#australia. The report is based on 500 metre pixel data on monthly time steps. Land use forest cover:

Results can be shown for the whole region (polygon), and separated by land use and forest cover classes which are likely to show different cover patterns and targets. Land use is divided into four broad classes: Conservation and natural environments, Agriculture, production native forests and plantation forests (no report), and other (no report). Agriculture is divided into grazing, crops and horticulture which are sub-divided into non-irrigated and irrigated. If forest is present land use is further divided into: non-forest, woodland forest and non-woodland forest. The area of each land use and forest class are shown as a map and chart. The report content is repeated for each land use and forest covers at least 1% of the area of the chosen region. Total vegetation Cover:

The total vegetation cover indicates where soil is likely to be protected from wind and or water hillslope erosion. Total vegetation cover for this month is shown on a map and chart classified into 4 classes.

• 71-100% High cover – protected from wind and usually water erosion (high rainfall, steep slopes, and erodible soils may need greater than 80, 90, 95 and up to 100% cover)

- 51-70% Moderate cover protected from wind erosion
- 31-50% Low cover not protected
- 0-30% Very Low cover not protected

Erosion protection: Wind erosion 50% total vegetation cover

The vegetation cover threshold required to prevent soil erosion is usually 50% to reduce wind erosion, 70% or 80% to reduce water (hillslope) erosion depending on the steepness and rainfall. Areas protected from erosion for the month:

- Map: water erosion protection (>70% cover) percentage area and hectares.
- Map: wind erosion protection (>50% cover) percentage area and hectares. Comparison with previous years:
 - Map: anomaly comparing this month to the average cover from the same month in previous years.
 - Map: deciles rank of month against the same month in previous years.

Anomalies and deciles until September 2019 are calculated comparing to the same months 2001 to 2019. Extra monthly data will be used to calculate anomalies and deciles post September 2019 as they become available. Time series monthly from January 2001 to current:

Erosion protection

- Wind erosion protection time series: percentage of the area of the region with greater than 50% cover for each month (orange lines). Horizontal lines are 10th (cover target) and 50th percentiles.
- Water erosion protection time series: percentage of the area of the region with greater than 70% cover for each month (blue line). Horizontal lines are 10th (cover target) and 50th percentiles.

Rainfall

• Millimetres rainfall each month (black line).

Each time series is also stacked by year. The black line shows the current year of data. Water erosion protection for higher rainfall and steeper slopes:

Water erosion protection on higher slopes. As slope increases, more cover is required to control water erosion. The thresholds reported are:

- the percentage area with pixels greater than 80% total cover.
- the percentage area with pixels greater than 90% total cover.
- the percentage area with pixels greater than 95% total cover.


Acknowledgment of data:

- 1. http://www.agriculture.gov.au/abares/aclump/land-use/alum-classification
- 2. http://www.agriculture.gov.au/abares/forestsaustralia/sofr/sofr-2018
- 3. https://www.dpi.nsw.gov.au/agriculture/pastures-and-rangelands/establishment-mgmt/production-management2/groundcover
- 4. MODIS Fractional cover algorithm:

https://doi.org/10.4225/08/5848a3f19a7b3



Grazing - Forest (non woodland)



Catchment Scale Land Use and Forests of Australia (2018) Derived from Catchment Scale Land Use of Australia (2018) and Forests of Australia (2018)

Anomaly show how many percetage points each

pixel is from

is, red pixels are about 20% lower than the

mean of that pixel. The mean

is only for the month of the map


using baseline

from 2001 to 2019.


the mean. That


Total Vegetation Cover [%]


Land use and forest cover




% Area protected from water erosion (>70%)








Proportion of vegetation cover class in area



% Area protected from wind erosion (>50%)

Total Vegetation Cover Anomaly [%]




Area not protected 0.0% of region (0 ha) È 73 Area protected 100.0% of region (157,325 ha) **Total Vegetation Cover Decile [%]** $\hat{\mathcal{S}}$ ୍ଚି Deciles show where the 1 N.1 2.3




record, from highest to lowest, for that month. That is, red pixels are in the lowest 10% of records for that month of the map using baseline from 2001 to 2019.

pixel value lies in the




Proportion of the area protected from wind erosion (Total Vegetation Cover > 50%)

Proportion of the area protected from water erosion (Total Vegetation Cover > 70%)



Grazing - Forest (non woodland) timeseries

Water erosion historical monthly area protected (Total Veg Cov>70%)



Wind erosion historical monthly area protected (Total Veg Cov >50%)